

The Incidence of Dwarf Novae in Large Area Transient Searches

A. Rau¹, R. Schwarz², S.R. Kulkarni¹, E.O. Ofek¹, M.M. Kasliwal¹, C. Brinkworth³, S.B. Cenko⁴, Y. Lipkin⁵ & A.M. Soderberg¹

¹ *Caltech Optical Observatories, MS 105-24, California Institute of Technology, Pasadena, CA 91125, USA*

² *Astrophysical Institute Potsdam, An der Sternwarte 16, 14482 Potsdam, Germany*

³ *Spitzer Science Center, California Institute of Technology, Pasadena, CA 91125, USA*

⁴ *Space Radiation Laboratory, California Institute of Technology, Pasadena, CA 91125, USA*

⁵ *School of Physics and Astronomy and Wise Observatory, Tel-Aviv University, Tel-Aviv 69978, Israel*

arne@astro.caltech.edu

ABSTRACT

Understanding and quantifying the contribution of known classes of transient and variable sources is an important lesson to be learned from the manifold of pre-cursors programs of the near-future large synoptic sky survey programs like SkyMapper, Pan-STARRS and LSST. With this goal in mind, we undertook photometric and spectroscopic follow-up observations of three recently reported unidentified transients. For two sources, WFI J132813.7–214237 and WFI J161953.3+031909, we show that unfortunate coincidences lead to their previous designation as transients. While the former is now interpreted as the spatial coincidence of a solar system object with faint background star, the latter is merely a cataclysmic variable unfortunately caught in and out of eclipse. The third candidate, ROTSE3 J160213.1–021311.7 is identified as an SU UMa-type dwarf novae with quiescent brightness of $R \sim 22.7$ and an outburst amplitude of ~ 5 mag. The fourth event, SDSS-SN15207, similarly shows evidence for a dwarf nova origin. Our main conclusion is that cataclysmic variables in their various avatars will contribute moderately to the population of transient objects.

Subject headings: binaries: eclipsing — stars: dwarf novae — novae, cataclysmic variables

1. Introduction

The study of transient and variable phenomena has historically been one of the main drivers of optical astronomy. This field is now widely expected to undergo a revival, thanks to the dawn of dedicated facilities, such as SkyMapper (Schmidt et al. 2005), the Panoramic Survey Telescope and Rapid Response System (Pan-STARRS; Kaiser et al. 2002) and the Large Synoptic Survey Telescope (LSST; Tyson 2005). A number of precursor surveys (e.g., Becker et al. 2004; Rykoff et al. 2005; Sako et al. 2005; Rau et al. 2006; Astier et al. 2006; Malacrino et al. 2006) are already informing us of the difficulty of finding genuine new classes of transients. Solar system objects (Malacrino & Atteia 2006) and flares from M dwarfs (Becker et al. 2004; Kulkarni & Rau 2006) are now demonstrated to be the most common variable objects.

In this paper we report on follow-up observations of four optical transients discovered in the ROTSE-III sky patrol (Rykoff et al. 2006), the MPG/ESO Wide Field Imager (WFI) search for orphan afterglows (Rau et al. 2006). and the SDSS supernovae search. Our findings show that cataclysmic variables (CVs) in general form an additional thin layer of foreground, which also must be pierced through before we can discover new classes of optical transients.

2. ROTSE3 J160213.1–021311.7

2.1. Discovery

The transient was discovered in ROTSE-III sky patrol fields on UT 2006 May 28.94 at an unfiltered magnitude of 18.2 ± 0.2 (Rykoff et al. 2006). No counterpart was found in previous ROTSE-III observations, the Digitized Sky Survey nor the 2-Micron All-Sky Survey. Following the discovery, the source brightened by ~ 1.4 mag within a day and quickly faded back to around 18th magnitude, where it seemed to stabilize for several days. Observations with the 1.5-m Russian-Turkish Telescope in Bakyrlytepe, Turkey, on June 1.88 UT, 2006, detected the source at $B = 18.96 \pm 0.02$, $V = 18.31 \pm 0.02$ and $R_c = 17.69 \pm 0.02$ (Rykoff et al. 2006). The $V - R$ color of 0.6 mag was commented to be redder than expected for an outburst of a cataclysmic variable. Thus, the source identification remained uncertain.

2.2. Photometry

Our photometric follow-up of the transient (Figure 1) started on UT 2006 June 4.95, two days after the discovery was announced and six days after the discovery itself. Our first epoch of observations were obtained with the 1-m telescope at the Wise Observatory, Israel, in the B , V , R and I bands. Monitoring in g and R with the automatic Palomar 60-inch telescope (P60) began two days later and a final epoch was obtained with the Large Field Camera at the Palomar 200-inch telescope (P200). A summary of the observations is presented in Table 1.

The first imaging epoch found the source at a brightness of $B = 19.83 \pm 0.12$, $V = 19.15 \pm 0.08$, $R = 18.70 \pm 0.05$ and $I = 18.81 \pm 0.08$. The subsequent observations with P60 traced the decay which was interrupted by several re-brightening phases with amplitudes of $\Delta M = 0.2 - 0.6$ mag and color changes from $g - R \sim 0.8$ mag to $g - R \sim 1.2$ mag during the variations (Figure 2). The total Galactic foreground extinction in the direction of the candidate is $E(B - V) = 0.276$ (Schlegel et al. 1998).

During the first nights of P60 observations the brightness of the transient allowed a variability study on short time scales using the individual consecutive 150 s R -band exposures (inset of Figure 2).

We inspected images generated by the Near-Earth Asteroid Tracking (NEAT) program¹ for past outburst episodes at the position of the transient. No source was detected in 17 epochs between June 1998 and July 2003 with limiting magnitudes (roughly R -band) of 20–21.5. Similarly, no X-ray counterpart was found in the ROSAT All-Sky survey² with a limit of 6×10^{-3} count s⁻¹ (0.1–2.4 keV). We conclude that the source is mostly in a quiescent state and rarely undergoes outbursts.

2.3. Spectroscopy

Spectroscopy was obtained with the Double-Beam Spectrograph (DBSP) mounted on the Hale-5-m telescope at Palomar Observatory (see Table 6). DBSP, as suggested by the name, operates with a dichroic and spectra can be obtained in two (blue and red) channels simultaneously. The spectrum (Figure 3) shows a flat continuum with strong emission features which we identify with $H\alpha$, $H\beta$ and $H\gamma$. The emission lines are clearly broadened

¹<http://skys.gsfc.nasa.gov/symorph/obs.html>

²<http://www.xray.mpe.mpg.de/cgi-bin/rosat/rosat-survey>

with a full width at half maximum ($45 \pm 2 \text{ \AA}$ for $H\alpha$) suggesting velocities of $> 2000 \text{ km s}^{-1}$. They are structured and double peaked in case of $H\beta$ and $H\gamma$. In addition, a strong HeII emission line at $\lambda 4686 \text{ \AA}$, blended with the Bowen fluorescence (Bowen 1934) of NIII, CIII and CIV is detected. The line properties are summarized in Table 3.

2.4. Identification of ROTSE3 J160213.1–021311.7

The decaying light curve matches a fast classical novae (CN) burst. However, the absolute peak magnitude of a fast CN is $\sim -8 \text{ mag}$ which would place the transient at a distance of 1 Mpc. The absence of a host galaxy makes this interpretation unlikely.

However, a dwarf nova hypothesis is entirely reasonable and in agreement with the emission spectrum. Dwarf novae constitute an important subgroup of cataclysmic variables which undergo outbursts with a brightness increase of 2–8 mag and repetition time scales of days to decades. The large amplitude ($> 4 \text{ mag}$, see Figure 2) of ROTSE3 J160213.1–021311.7 along with the outburst duration of $> 30 \text{ days}$ resembles a *superoutburst* of SU UMa-like DNe. Superoutburst light curves of these sources typically exhibit a plateau phase with a slow decay of $\sim 1 \text{ mag}$ in 8–10 days after the initial brightening, followed by a steep decline with periodic luminosity modulations, called *superhumps*. The observed light curve differs by showing an early flare and a shallower (quasi-exponential) decay at late times. However, we found that this temporal behavior is very similar to the superoutburst of the SU UMa dwarf nova RU Peg in 1981 (la Dous et al. 1985). Although our data are too sparse to detect clear evidence for superhumps, the re-brightenings and variability on short times scales (see inset of Fig. 2) could be interpreted as signatures of such.

The spectrum shows strong similarities to IY UMa (an SU UMa DN) during its superoutburst in 2000 (Rolfe et al. 2005). The large flux ratio of $\text{HeII}/H\beta \sim 1.2$ is also a popular indicator for a highly magnetic CV, e.g. a polar or an intermediate polar. However, the temporal behavior of ROTSE3 J160213.1–021311.7 with its large amplitude outburst from a long lasting quiescent state is entirely inconsistent with the long-term variability expected for a magnetic CV. In addition, the double peaked emission lines and lack of pronounced cyclotron lines clearly provide further evidence that this system is a typically disk CV.

Finally, we note the large $g - R$ color (~ 1) could suggest a cold accretion disk in the system resulting from a low mass accretion rate. A low mass transfer is reminiscent of cataclysmic variables in the period gap between 2 and 3 hours (e.g., Robinson 1983; Ritter & Kolb 1995), which extends to $\sim 4 \text{ hrs}$ for dwarf novae (Shafter et al. 1986). Thus, we conjecture that ROTSE3 J160213.1–021311.7 is a hibernating (Shara et al. 1986) SU UMa

dwarf nova.

3. WFI J161953.3+031909

3.1. Discovery

In 1999, a transient and variability survey (dedicated to the search for orphan afterglows) was performed at the MPG/ESO 2.2-m telescope equipped with the Wide Field Imager (WFI). 12 square degrees were monitored in up to 25 nights down to a limiting magnitude of $R = 23$ (Rau et al. 2006). The search netted four candidates after filtering for moving or known solar system objects and known variables. For two sources, a classification as a CV and a flare star were suggested based on their temporal behavior. Follow-up observations for remaining two will be described here and in the following section.

WFI J161953.3+031909 is an uncataloged transient found in the WFI survey when it showed a brightening from a persistent $R = 19.9$ to $R = 17.5$ within two days at UT 1999 June 17.14. Based on the subsequent optical decay and the detection of a faint X-ray source in the ROSAT all-sky survey, the source was preliminarily classified as a dwarf nova of the SU UMa class.

3.2. Photometry

Photometry was obtained in seven epochs in May 2006 with the P60. A total of 220 R -band images with exposures of 150 s were taken. The photometric calibration was tied to the original WFI discovery image (see Rau et al. 2006, for details). Relative photometry was applied to correct the series of P60 observations to a common reference system. The light curve revealed periodic occultations (Figure 4) and we derived a firm ephemeris from the timing of six individual eclipses as

$$\text{HMJD}(T_{\text{ecl}}) = 53880.2503(12) + E \times 0.099401(9). \quad (1)$$

with the numbers in brackets giving the uncertainty in the last digits. The accumulated cycle count between the WFI and P60 observations is too large to derive a unique period solution from 1999 to 2006.

3.3. Spectroscopy

Spectroscopy was performed with the Low Resolution Imager and Spectrograph (LRIS; Oke et al. 1995) mounted on the Keck I 10-m telescope (see Table. 6). The LRIS light beam is split by a dichroic and spectra were obtained in the Blue and Red channel simultaneously. We obtained two 900 s spectra with mid-exposure times corresponding to orbital phases of ~ 0.36 and ~ 0.49 . The spectra show blue continua with a number of strong emission features which we identify with Balmer emission ($H\alpha \dots H\gamma$) and HeI at $\lambda 5876$ and $\lambda 6678$. All lines are double-peaked and clearly broadened with respect to the instrumental line width (see inset of Figure. 5). In addition, a broad emission feature centered around $\lambda = 4665 \text{ \AA}$ is found at phase 0.49. We associate this with a Bowen blend of NIII, CIII and CIV. A summary of the line properties is given in Table 4.

3.4. Identification

The periodic light curve and the spectrum resemble those of a dwarf nova in quiescence. The detection of deep orbital eclipses (Figure 4), together with the double peaked emission lines indicate a high inclination of the system. Out of eclipse, the light curve appears flatter than typically observed in eclipsing dwarf novae. In addition, no strong asymmetry is seen between ingress and egress, suggesting that the hot spot, the region where the gas stream collides with the accretion disk, would be weaker than usual. This is in agreement with the low mass transfer rate assumed for CVs in the period gap, in which WFI J161953.3+031909 falls.

The double peaked emission lines (Figure 5), the absence of pronounced cyclotron features and the non-detection of HeII indicate a disk system. The detection of the Bowen blend around $\lambda 4665$ bears evidence of irradiation in the disk or the secondary. HeII emission, which leads to the Bowen fluorescence, is not detected, though. This suggests a considerable HeII opacity.

The orbital period of ~ 143 min implies that observations taken at the same UT but over different nights, will result in the source being observed at nearly identical orbital phases. In retrospect, we now realize that the original survey data were obtained at phases at which the object exhibited extreme brightness. The maximum brightness in the WFI detections of $R = 17.5$ corresponds to the median brightness out of eclipse (see Figure. 4) and the (falsely interpreted) quiescent brightness of $R = 19.9$ is similar to what is observed during the eclipse.

Summarizing, we found no evidence for a dwarf nova outburst of WFI J161953.3+031909

and thus are left with the rough classification of the source as a disk CV. Indeed, the brighter quiescent magnitude than previously thought, implies $L_X/L_{opt} \sim 0.1$, consistent with a larger variety of CVs.

4. WFI J132813.7–214237

4.1. Discovery

The source was detected on June 26.07 UT, 1999, at a maximum brightness of $R = 19.9$. As a result of the scheduling, no observations could be obtained during a subsequent decay. However, a faint uncataloged and persistent object ($R = 21.3$) was found, offset by $0''.8$ from the location of the transient source. This apparent association allowed for a possible extragalactic origin of the transient with the persistent source being the underlying host galaxy. It was therefore interpreted as the best candidate for an orphan afterglow found in the survey. However, given the sparse data coverage, other interpretations (extra-galactic, galactic and even solar system) could not be excluded.

4.2. Spectroscopy

Spectroscopy of the of the persistent source associated with WFI J132813.7–214237 was performed with LRIS. We find strong absorption features which we identify with Ca H&K, g-band, $H\beta$ and $H\alpha$ (Fig. 6). The narrow lines allow to estimate the velocity with respect to the local standard rest frame to $0 \pm 50 \text{ km s}^{-1}$. No significant emission lines are detected.

4.3. Identification

The spectrum is consistent with that of a galactic G-star. Thus, we can rule out an extra-galactic origin for the persistent source. As a result, there is little justification to assume that the transient source was extra-galactic. Instead the detection was likely the misleading alliance of a moving solar system object superimposed on a background star. The position in the ecliptic plane (ecliptic coordinates $\lambda = 208.5^\circ$ & $\beta = -11.6^\circ$) supports this scenario.

5. SDSS-SN15207

5.1. Discovery

SDSS-SN15207 was discovered on 2006 October 11 as part of the Sloan Digital Sky Survey (SDSS) search for supernovae at RA(J2000)=20^h58^m43^s45, Dec(J2000)=01°02′47″0. SDSS imaging prior to the eruption shows no underlying host galaxy to a limit of $r=24$. The SDSS optical light curve³ showed a peak magnitude of $r=19.8$ on Oct 12 followed by a decline by 1.2 mag over 5 days. Continued monitoring with P60 during the first 80 days did not detect additional outbursts to a limiting magnitude of $r=22$.

5.2. Spectroscopy

We obtained a spectrum with the Gemini Multi Object Spectrograph (GMOS) on Gemini-South on 2006 November 17 (Table 6). At this time, the event had faded to $r \sim 23$. This resulting low-signal-to-noise spectrum (Figure 7) shows a blue continuum and a single emission line (FWHM= 1000 ± 200 km/s), which we identify with H α . The line centroid suggest a Galactic origin.

We also obtained a 4.77 ks exposure with the X-ray telescope aboard Swift on 2006 November 4. No counterpart was found with a 2σ flux limit of $> 6 \times 10^{-14}$ ergs cm⁻² s⁻¹ (0.2–10 keV).

5.3. Identification

The peak magnitude of this transient rules out the possibility that it is a nova in the Milky Way. The optical light curve, in particular, the steep decline of 1.2 mag in 5 days and 1.6 mag in 9 days does not suggest a typical supernova either. On the other hand, the observations are consistent with a super outburst (> 4 mag) of a Galactic dwarf nova. The non-detection of the quiescent counterpart to $r = 24$ and the lack of significant foreground extinction in the optical colors and spectrum, suggest SDSS-SB15207 to have been located at a relative large distance from the earth. Assuming an absolute quiescence magnitude of $R = 12$ (white dwarf with $T = 11000$ K, $M = 0.6 M_{\odot}$; Bergeron et al. 1995), the lower limit on the distance is 2.5 kpc.

³<http://sdssdp47.fnal.gov/sdsssn/candidates/examineCand.php?cid=15207>

6. Discussion and Conclusion

Here we reported on photometric and spectroscopic follow-up observations for four recently communicated transient sources. We found that WFI J132813.7–214237, the best orphan afterglow candidate detected in the WFI survey (Rau et al. 2006), was with high probability a misidentified solar system object spatially coincident with a background source. Next, the potential outburst light curve of a second source from the same survey, WFI J161953.3+031909, was recognized as serendipitous observations in and out of eclipse of a high-inclination cataclysmic variable. Finally, ROTSE3 J160213.1–021311.7, discovered by ROTSE-III (Rykoff et al. 2006) and (potentially SDSS-SN15207) were identified as the superoutbursts of SU UMa-like dwarf nova.

In the following, we restrain from exploring the two WFI sources any further, except noting again the interesting coincidences which had lured us to incorrect conclusions previously. Instead, we want to address the implications of the faint quiescent magnitudes of ROTSE3 J160213.1–021311.7 ($g \sim 23$) and SDSS-SN15207 ($r > 24$). We start by noting that a very similar event (1955+22C VAR VUL 05 ⁴) has been discovered in August 2005 with 15.8 mag at maximum and an amplitude of at least 9 mag.

This quiescent faintness of these three events indicates an important factor for future wide-field transient searches, namely, dwarf novae which stay most of the time below the limiting magnitude of a survey, but appear as new transient sources during outbursts. This includes CVs which appear faint in quiescence because of large distance or Galactic extinction and intrinsically low-magnitude systems. The latter group is dominated by systems with low mass transfer, typical for CVs in the period gap and in hibernation (Shara et al. 1986). As a matter of fact, the vast majority of non-magnetic CVs with $P_{orb} < 3$ hrs are dwarf novae ($\sim 83\%$) while this is true only for $\sim 36\%$ with $P_{orb} > 3$ hrs (Aungwerojwit et al. 2006).

As an example, let us consider the proposed 3π steradian survey with PanSTARRS-1 (PS1) which is expected to reach a limiting magnitude of $R \sim 23.2$ and $R = 24.9$ in a single 30s and combined images over 3 yrs, respectively. In such a survey, sources similar to ROTSE3 J160213.1–021311.7 at twice the distance would no longer be detectable in quiescence, but only during the rare outburst periods. This directly leads to the question of how important these events will be in future wide-field surveys.

The expected number of DNe events in an all-sky snapshot can be obtained rather crudely, given the large number of involved parameters (e.g., intrinsic quiescent magnitude, superoutburst amplitude, superoutburst cycle). Let the quiescent absolute magnitude of the

⁴AAVSO Alert Notice 325

DN be $M_R \sim 12$ (Bergeron et al. 1995). For the volume distribution we assume a local scale height of 300 pc perpendicular to the Galactic disk and isotropy in the radial direction in the disk. Furthermore, we consider for simplicity an upper limit for the distance in the Galactic Plane of 2 kpc, above which extinction will dominate at low Galactic latitude. For the observed local density⁵ of dwarf novae we use $\rho \sim 3 \times 10^{-5} \text{ pc}^{-3}$ (Schwope et al. 2002) and theory allows for even higher values (10^{-4} to 10^{-3} pc^{-3} ; Kolb 1993; de Kool & Ritter 1993). The resulting number of expected dwarf novae in the considered volume, independent of their activity, is given in Table 5 for three different ranges of Galactic latitude.

How many of those will be in superoutburst at any given time? Assuming a mean cycle duration of 1 yr and a mean plateau length of 10 days we find that between 40 and 1.5×10^3 events, depending on the Galactic latitude and assumed space density, can be expected in an all-sky snapshot with maximum depth. Most of these ($\sim 86\%$) will be at low and medium Galactic latitude and only a small fraction ($\sim 14\%$) is expected to occur at $|b| > 45^\circ$.

The space density of CVs decreases rapidly with distance from the Galactic Plane. Similarly, at low Galactic latitude extinction limits the observable. As a result, the number of dwarf novae with quiescent brightness below the survey limiting magnitude (here $R \sim 25$) will be negligible. However, shallower surveys will detect less quiescent systems as indicated in Figure 8. Thus, mainly these surveys will be contaminated by dwarf novae superoutbursts with undetected quiescent counterparts.

The yearly all-sky rate of less than 4×10^4 detectable superoutbursts is in the range of the local volume rates ($\text{Gpc}^{-3} \text{ yr}^{-1}$) for core-collapse SNe ($\sim 5 \times 10^4$; Cappellaro et al. 1999) but considerably lower than the rates of novae ($\sim 10^8$; Kulkarni & Rau 2006) and Galactic M-dwarf flares (10^8 yr^{-1} ; Kulkarni & Rau 2006). In addition, one has to remember that the theoretical CV space densities typically imply a large fraction of systems with cycle periods of years to decades (e.g., WZ Sge). Thus, the yearly all sky rate is probably much less than value given above. This suggests that dwarf nova superoutbursts will not produce a dominant contamination in future wide-field surveys. However, they will still be of concern for observers mining the optical transient sky.

How can the small number of dwarf novae outbursts be distinguished in practice from other kinds of interesting transients? The most important factors will be the choice of cadence and the field selection. A good sign for a DN will be a location close to the Galactic disk or inner Galaxy. Indeed, the positions of ROTSE3 J160213.1–021311.7 ($l = 8^\circ$, $b = 35^\circ$) and also of 1955+22C VAR VUL 05 ($l = 60^\circ$, $b = -4^\circ$) are consistent with being part of

⁵Note that the observationally derived space densities are uncertain by at least a factor of ten in both directions. A detailed assessment of possible selection effects in surveys can be found in Patterson 1984.

the disk population. However, the Galactic Plane is known to harbor plentiful of variable and transient sources. Choosing the right observing strategy (cadence, filter) will help to distinguish between short lived phenomena (e.g., flare stars) and dwarf novae. The main confusion at high Galactic latitude will be by novae and supernovae. Novae, however, will be distinguishable by their enormous brightness (if Galactic) or by their association with a nearby galaxy. Supernovae may similarly identified by their host galaxies. Spectroscopy of the remaining candidates would require significant amounts of time on large telescopes and is thus not a feasible approach. Photometric colors vary considerably from one event to another and will therefore not be useable as reliable indicators either. The best confirmation, however, can be obtained by photometric follow-up searches for the expected superhump structures.

We close this paper noting the role of future large surveys for our general understanding of cataclysmic variables. Complete wide-area surveys on long times scales will allow the in-depth study of the distribution of orbital periods and outburst behaviors (amplitudes, cycles). This will provide important informations on the evolution of CVs and the processes of mass accretion in these systems.

We acknowledge helpful discussions with J. Greiner, E. Mason and L. Bildsten. We thank M. Sako and A. Becker for the timely alert about the eruption of SDSS-SN15207. Furthermore, we thank the anonymous referee for constructive comments. This work is supported in part by grants from the National Science Foundation and NASA.

REFERENCES

- Astier, P., Guy, J., Regnault, N., 2006, *A&A*, 447, 31
- Aungwerojwit, A., Gänsicke, B.T., Rodriguez-Gil, R., et al., 2006, *A&A*, 455, 659
- Becker, A.C., Wittman, D.M., Boeshaar, P.C., et al., 2004, *ApJ*, 611, 418
- Bergeron, P., Wesemael, F., & Beauchamp, A., 1995, *PASP*, 107, 1047
- Bowen, I.S., 1934, *Pub. A.S.P.*, 46, 146
- Cappellaro, E., Evans, R., & Turatto, M., 1999, *A&A*, 351, 459
- de Kool & Ritter, H., 1993, *A&A*, 267, 397
- Hellier, C., 2001, in *Cataclysmic Variable Stars*, Springer

- Kaiser, N., et al., 2002, *Proc. SPIE*, 4836, 154
- Kolb, U., 1993, *A&A*, 271, 149
- Kulkarni, S.R., & Rau, A., 2006, *ApJ*, 644, L63
- Malacrino, F., & Atteia J.-L., 2006, *GCN*, 4705
- Malacrino, F., Atteia, J.-L., Boer, M., et al., 2006, *A&A*, 459, 465
- la Dous, C., Verbunt, F., Schoembs, R., 1985, *MNRAS*, 212, 231
- Landolt, A.U., 1992, *AJ*, 104, 340
- Oke, J.B., et al., 1995, *PASP*, 107, 375
- Patterson, J., 1984, *ApJS*, 54, 443
- Rau, A., Greiner, J., & Schwarz, R., 2006, *A&A*, 449, 79
- Ritter, H., Kolb, U., 1995, in *X-ray binaries*, ed. W.H.G. Lewin, J. van Paradijs, E.P.J. van den Heuvel (Cambridge: Cambridge University Press), 578
- Robinson, E.L., 1983, in *Cataclysmic Variables and Related Objects*, ed. M. Livio & G. Shaviv (Dordrecht:Reidel), 1
- Rolfe, D.J., Haswell, C.A., Abbott, T.M.C., 2005, *MNRAS*, 357, 69
- Rykoff, E., Aharonian, F., Akerlof, C.W., et al., 2005, *ApJ*, 631, 1032
- Rykoff, E.S., Aretakis, J. & Guver, T., 2006, *ATel*, 824
- Sako, M., Romani, R., Frieman, J., et al., 2005, *astro-ph/0504455*
- Schlegel, D.J., Finkbeiner, D.P., & Davis, M., 1998, *ApJ* 500, 525
- Schmidt, B. P., Keller, S. C., Francis, P. J., Bessell, M. S., 2005, *AAS*, 206, 1509
- Schwope, A., Brunner, H., Buckley, D., et al., 2002, *A&A*, 396, 895
- Shafter, A.W., Wheeler, J.C., & Cannizzo, J.K., 1986, *ApJ*, 305, 261
- Shara, M.M., Livio, M., Moffatt, A.F.J., & Orio, M., 1986, *ApJ*, 311, 163
- Tyson, A., 2005, in *ASP Conf. Ser. 339: Observing Dark Energy* (eds Wolff, S.C., & Lauer, T.R., 95-107




Fig. 1.— P60 image of the field around ROTSE3 J160213.1–021311.7 taken on June 7, 2006. Astrometry was performed based on USNO-B1.0 positions of ten stars in the neighborhood of the transient. The coordinates of the transient (circle) are $\alpha_{J2000}=16:02:13.06$ & $\delta_{J2000}=-02:13:12.4$. A comparison star with $R = 16$ mag located $0''.9$ West and $75''.6$ South at $\alpha_{J2000}=16:02:13:00$ & $\delta_{J2000}=-02:14:28.0$ is marked by the horizontal dash.




Fig. 2.— P60 light curve of ROTSE3 J160213.1–021311.7 . Photometric calibrations (Vega system) were obtained using observations of the Landolt standard star SA108-475 (Landolt 1992). A clear monotonic brightening by ~ 0.8 mag over about 30 min was observed on June 7 (inset). Variability was also detected on June 8–10 at lower significance. The epoch of our DBSP spectroscopy is indicated by the vertical dash.




Fig. 3.— DBSP spectrum of ROTSE3 J160213.1–021311.7 . The prominent Balmer and HeII emission features (labeled) are structured and a double peaked shape is visible for $H\beta$ (inset) and $H\gamma$. The inset shows also the region around the HeII line at 4686 $\text{\AA}$ revealing a strong Bowen blend of CIII, CIV and NIII.




Fig. 4.— Phase-folded P60 light curves of WFI J161953.3+031909 using the ephemeris given in Eq. 1 shown twice for clarity. Data taken on May 25th (points) and 26th (empty lozenge) reveal alterations of the out-of-eclipse variability, as well as a changes of the overall brightness level of up to 0.5 mag. Additional observations were performed on May 12, 13, 15, 16, & 17, 2006 (see electronic supplement). Horizontal dashes mark the phases of the Keck/LRIS spectroscopy observations (see Figure 5)




Fig. 5.— LRIS spectra of WFI J161953.3+031909 taken 2006 April 22 at orbital phases ~ 0.36 (bottom, grey) and ~ 0.49 (top). Prominent Balmer and HeI emission lines are indicated. The emission lines show a double peaked structure, which is shown for $H\alpha$ (phase ~ 0.49) in the inset. In addition, a prominent Bowen blend is seen at phase ~ 0.49 shortwards of $H\beta$.




Fig. 6.— LRIS spectrum of the quiescent counterpart to WFI J132813.7–214237 taken 2006 April 23. Prominent absorption features are indicated.




Fig. 7.— Gemini GMOS spectrum of SDSS-SN15207 obtained 2006 November 17, 36 days after the discovery.




Fig. 8.— Predictions for the number of dwarf novae which can be detected in a survey with a given limiting R -band magnitude. The solid, dashed and dotted lines show the expectations for three ranges of Galactic latitude. Note, that the sharp turnover for $|b| < 10^\circ$ is caused by the distance limit in the Galactic Plane (2 kpc) which was assumed to account for foreground extinction. At $|b| > 45^\circ$ nearly all quiescent system will be detected above $R = 22$.

Table 1: ROTSE3 J160213.1–021311.7 imaging log (for electronic version).

MJD	Telescope	Filter	Exposure (s)	Brightness (mag)
53890.94	Wise	B	840	19.83±0.12
53890.94	Wise	V	600	19.15±0.08
53890.93	Wise	R	600	18.70±0.05
53890.96	Wise	I	600	18.81±0.08
53893.19	P60	g	1350	19.16±0.08
53893.19	P60	R	1350	18.42±0.05
53894.21	P60	g	1350	20.01±0.09
53894.21	P60	R	1350	18.79±0.05
53895.22	P60	g	1350	20.01±0.11
53895.22	P60	R	1350	19.22±0.07
53896.26	P60	g	1350	19.86±0.15
53896.26	P60	R	1350	19.05±0.09
53898.21	P60	g	1350	20.82±0.29
53898.21	P60	R	1350	20.34±0.16
53899.25	P60	g	1350	20.84±0.22
53899.25	P60	R	1350	19.78±0.11
53906.16	P60	g	1350	22.26±0.21
53906.16	P60	R	1350	21.29±0.22
53908.20	P60	R	1350	21.43±0.19
53914.29	P60	g	1350	23.10±0.40
53914.29	P60	R	1200	21.87±0.29
53929.19	P60	R	2700	22.53±0.37
53949.19	P60	R	2700	22.18±0.28
53968.23	P200	g'	600	23.5±0.2
53968.24	P200	r'	600	22.7±0.1

Note. — All magnitudes are given in the Vega sytem.

Table 2: Log of the spectroscopic observations.

	Instrument	Setup	Exposure (s)
ROTSE3 J160213.1–021311.7:			
2006 June 5	Hale-5m/DBSP	300/3990	2100
	Hale-5m/DBSP	316/7500	2100
WFI J161953.3+031909:			
2006 April 22	Keck/LRIS	400/3400	2×900
	Keck/LRIS	400/8500 (7608 Å)	2×900
WFI J132813.7–214237:			
2006 April 23	Keck/LRIS	400/3400	3600
	Keck/LRIS	400/8500 (7608 Å)	3600
SDSS-SN15207:			
2006 November 17	Gemini-S/GMOS	150/7170	2×1750

Note. — In column three we list the grism/grating (number of grooves/blaze angle) and (for the red DBSP and LRIS gratings) the central wavelength.

Table 3: ROTSE3 J160213.1–021311.7 emission line characteristics.

Species	Flux $\times 10^{-15}$ erg/cm ² /s	–EW (Å)	FWHM (Å)
H γ λ 4340	1.6±0.3	19±4	42±7
HeII λ 4686	1.9±0.3	25±5	42±5
H β λ 4861	1.7±0.2	22±5	41±7
H α λ 6563	2.6±0.2	58±5	45±2

Table 4: WFI J161953.3+031909 emission line characteristics.

Species	Phase ~ 0.36		Phase ~ 0.49	
	Flux $\times 10^{-15}$ erg/cm ² /s	-EW (Å)	Flux $\times 10^{-15}$ erg/cm ² /s	-EW (Å)
H δ λ 4102	<1.2	<5	1.4 ± 0.2	-3 ± 1
H γ λ 4340	1.5 ± 0.5	6 ± 2	2.7 ± 0.2	4 ± 1
Bowen λ 4665	<2.2	<9	5.5 ± 0.5	15 ± 5
H β λ 4861	2.8 ± 0.3	13 ± 1	3.7 ± 0.2	13 ± 1
HeI λ 5876	1.0 ± 0.2	7 ± 1	1.5 ± 0.4	6 ± 1
H α λ 6563	3.9 ± 0.3	34 ± 3	5.4 ± 0.3	34 ± 3
HeI λ 6678	0.4 ± 0.1	3.4 ± 0.4	0.6 ± 0.1	3.6 ± 0.5

Table 5: Dwarf novae number counts and outburst rates.

Galactic latitude	N_{DN}^1		N_{SO}^2	
	$\rho = 3 \times 10^{-5}$ pc ⁻³	$\rho = 10^{-3}$ pc ⁻³	$\rho = 3 \times 10^{-5}$ pc ⁻³	$\rho = 10^{-3}$ pc ⁻³
$ b < 10^\circ$	$\sim 1.7 \times 10^3$	$\sim 5 \times 10^4$	~ 40	$\sim 1.5 \times 10^3$
$10^\circ < b < 45^\circ$	$\sim 1.8 \times 10^3$	$\sim 5 \times 10^4$	~ 40	$\sim 1.5 \times 10^3$
$ b > 45^\circ$	$\sim 5 \times 10^2$	$\sim 1.5 \times 10^4$	~ 15	$\sim 4 \times 10^2$

Note. — ¹: total number of dwarf novae within the considered volume. See text for details. ²: number of superoutbursts in an all-sky snap shot. Note, that the theoretical DN density includes a large population of events with superoutburst cycles of decades. Thus, the presented numbers are optimistic. Results will change if a different scale height (e.g., 150 pc; Patterson 1984) is adopted.